

Perianal surgery

There may be operations for external hemorrhoids, skin folds after hemorrhoids or abscesses. Before the operation, you will receive an injection of local anesthetic medication at the site in question, so that the operation is painless. The anesthesia lasts for 1-3 hours.

Preparation

If you are taking blood-thinning medicines (eg Heart Magnyl , Marevan , Eliquis , Xarelto or Clopidogrel) or have a pacemaker, inform your clinic well in advance of the examination.

Treatment

Performed under local anesthesia, which is applied via plugs in or around the hemorrhoids.

After treatment

The first few days there will often be some pain around the rectal opening. It will therefore be a good idea to take 2 tabl. Pinex 500mg and 2 tabl. Ipren at 200 mg 3 x daily.

The wounds after the operation are always left open and they heal spontaneously within 3-6 weeks.

It can fluid from the wounds until they are healed.

It may bleed from the wounds for the first 1-2 days and then during bowel movements, especially if it is hard.

The stool

It is important that the stool is kept going and does not stop.

It is important to emphasize that future intake of plant fiber will in many cases prevent the re-emergence of the disease.

Complications

Very rarely seen, bleeding and infection may occur. In case of signs of major bleeding or signs of inflammation, contact the clinic, emergency doctor or emergency room immediately. Very rarely, symptoms of a tear and impaired sphincter function may occur.